

A NEW COLOR FORM OF CORNEL FROM NEW JERSEY

Fred W. Oswald

*CORNUS RACEMOSA* f. *CAERULEOCARPA* Oswald, f. nov.

Haec forma a forma typica speciei fructibus atro-caeruleis recedit.

This form differs from the typical form of the species in



having deep-blue fruit instead of the normal white fruit.

The type of this form was collected by myself at the Alpine Scout Camp, Alpine, Bergen County, New Jersey, on August 30, 1955, and is deposited in the H. N. Moldenke Herbarium at Yon-

kers, New York. The accompanying plate was drawn from fruiting material taken from the type plant on September 12, 1955.

-----

NOTES ON NEW AND NOTEWORTHY PLANTS. XXI

Harold N. Moldenke

AZALEA ALBICANS (Waterer) Moldenke, comb. nov.

Rhododendron albicans (Waterer) Zabel, Mitt. Deutsch. Dendr. Gesell. 11: 30. 1902.

AZALEA KOSTERIANA (C. K. Schneid.) Moldenke, comb. nov.

Rhododendron kosterianum C. K. Schneid., Ill. Handb. Laubholz. 2: 499. 1911.

AZALEA OBTUSA f. JAPONICA (Maxim.) Moldenke, comb. nov.

Rhododendron obtusum f. japonicum (Maxim.) Wils. in Wils. & Rehd., Mon. Azal. 33. 1921.

x ERIOCAULON HESSII Moldenke, nom. nov.

Eriocaulon angustisepalum x mutatum H. Hess, Bericht. Schweiz. Bot. Gesell. 65: 173--174. 1955.

ERIOCAULON JAEGERI Moldenke, sp. nov.

Herba parva annua acaulescens; foliis parvis caespitosis tenuissime membranaceis vel translucentibus fenestratis graminoidibus utrinque glabris subulato-acutis; pedunculis paucis filiformibus rectis stramineis tricostatis nitidis glabris; vaginis arcte adpressis tenuissime membranaceis glabris oblique fissis; capitulis hemisphaericis albis; bracteis receptaculi late obovatis nigris ad apicem late rotundatis apiculatis, ad basin cinea-tis, glabris nitidis.

Small, annual, stemless herb; leaves few, tufted, very thin-membranous or translucent, fenestrate, very pale-green, linear and grass-like, about 3.5 cm. long and 1.5--2 mm. wide at and below the middle, glabrous on both surfaces, subulate-acute at the apex; peduncles few, 3--6 per plant, filiform, erect, 3--13.5 cm. long, 3-costate, stramineous, shiny, glabrous; sheaths slender, closely appressed, 4--4.5 cm. long, very thin-membranous, glabrous, obliquely split at the apex, the blade 5--7 mm. long, erect, appressed, not spreading, glabrous, the tip long-attenuate or subulate; heads hemispheric, 4--9 mm. wide, white; receptacular bractlets broadly obovate, black, concave, about 1.6 mm. long and wide, broadly rounded and apiculate at the apex, cuneate to the base, glabrous, shiny; staminate florets: sepals 3, blackish, oblong-linear, concave, 1.2--1.4 mm. long, about 0.2 mm. wide, acute at the apex, glabrous; petals 3, united into a slender white tube about 1.6 mm. long, the free tips unequal